



Gallovac 9R[®]

Lyophilisate suspension for injection live attenuated
Salmonella gallinarum strain 9R vaccine

STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS:

Each (0.2 ml) contains:

- Active substance:
Each dose contains a minimum of 2.0×10^7 CFU of freeze-dried live attenuated *Salmonella gallinarum* vaccine prepared from the reference strain 9R.
- Other ingredients:
Bovine serum, sucrose and Sodium glutamate.

INDICATIONS:

For active immunization of layers and breeders chickens against *Salmonella gallinarum* (fowl typhoid) / pullorum and partial control of *Salmonella Enteritidis* infections.

CONTRAINDICATIONS:

Do not vaccinate sick or stressed birds.

ADVERSE REACTIONS:

Very rare; small nodular reactions may occur at the sight of inoculation which disappear later.
If you notice any side effects please inform your veterinarian.

TARGET SPECIES:

Chickens.

DOSAGE:

- Dosage is 0.2 ml per bird subcutaneously into the lower part in the back of the neck.
- Primary vaccination: at 6-8 weeks of age.
- Secondary vaccination: at 14-16 weeks of age.

ADVICE ON CORRECT

ADMINISTRATION:

- Apply usual aseptic procedures.
- Use only clean, antiseptic or disinfectant free materials for the preparation of the vaccine solution.
- Dissolve the freeze-dried pellet vaccine with cold and sterile solvent provided.
- Use 100 ml diluent for each 500 doses vaccine vial and 200 ml solvent for 1000 doses vaccine vial.

- Add approximately 3 ml of the solvent to the freeze-dried vaccine aseptically using a sterile syringe and mix well. Withdraw the dissolved vaccine into the syringe and add it to the solvent bottle.
- Use the vaccine immediately after reconstitution (within two hours in cool conditions).
- Exposure to infection should be avoided for 14 days following vaccination to allow immunity to be developed.

WITHDRAWAL PERIOD:

Zero days.

SPECIAL STORAGE PRECAUTIONS:

- Store in a refrigerator (2°C-8°C).
- Do not freeze.
- Protect from direct sunlight.
- Keep out of the reach and sight of children.
- The vaccine remains stable for 18 months from production date stated on the label under the recommended storage conditions.
- Shelf life after reconstitution: Use the vaccine within 2 hours after reconstitution.

SPECIAL WARNING(S):

special warnings for each target species:
Vaccinate healthy birds only.

Special precautions for use in animals:

Not applicable.

Special precautions to be taken by the person administering the medicinal product to animals:

Although Gallovac 9R vaccine is not pathogenic for humans; care should be taken to avoid accidental injection. Should accidental injection occur immediate medical care should be sought.

Use during lay, fertility:

Do not vaccinate layers and breeders before two weeks and during laying period. can be administered during the laying period in high risk-areas.

Interaction with other medicinal products and other forms of interaction:

Do not use antibiotics or other antibacterial substances for 7 days before vaccination.

Overdose:

10 X of the dose of a vaccine:

Rare: small nodular reactions may occur at the sight of inoculation which disappear later.

If you notice any side effects please inform your veterinarian.

SPECIAL PRECAUTIONS FOR DISPOSAL:

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

OTHER INFORMATION:

Gallovac 9R is prepared from *Salmonella gallinarum* strain 9R for active immunization of chickens against *Salmonella gallinarum* (fowl typhoid) / pullorum and partial control of *Salmonella Enteritidis* infections.

Package sizes:

Vials of: 500 doses 1000 doses.

FOR VETERINARY USE ONLY.



Jovac

MANUFACTURING AUTHORIZATION HOLDER AND MANUFACTURED BY:

JORDAN BIO INDUSTRIES CENTER
Tel. +962 6 5232162, Fax: +962 6 5232210
P.O. Box 43, Amman 11941- Jordan
E-mail: info@jovaccenter.com, www.jovaccenter.com

9RINE022.04
12/05/2022