



Jovaplastm C®

Inactivated suspension for injection vaccine
against Contagious Caprine Pleuropneumonia

STATEMENT OF THE ACTIVE SUBSTANCE(S) AND OTHER INGREDIENTS:

- Active substance:
Each dose (1 ml) contains a minimum of 0.15 mg of *Mycoplasma capricolum* subspecies *capripneumoniae* (Mccp) formally known as strain F38.
- Other ingredients:
Saponin.

INDICATIONS:

Jovaplastm C is indicated for the vaccination of goats at 3 months of age or older for protection against Contagious Caprine Pleuropneumonia.

CONTRAINDICATIONS:

Do not vaccinate sick or stressed animals.

ADVERSE REACTIONS:

Very Rare: A very small lump at the site of injection may appear in vaccinated animals. This lump usually disappears within 48 hours. If you notice any side effects please inform your veterinarian.

TARGET SPECIES:

Goats.

DOSAGE:

- Administer 1 ml subcutaneously, (preferably in the elbow region) at 3 months of age or older.
- Re-vaccinate with one recommended dose (1 ml) every 6 months.
- Vaccination of healthy goats in endemic and high risk areas is highly recommended.
- Routine vaccination of goats over 3 months of age.

ADVICE ON CORRECT ADMINISTRATION:

- Shake well before use.
- Apply usual aseptic procedures.
- Avoid stress when vaccinating.

WITHDRAWAL PERIOD:

Zero days.

SPECIAL STORAGE PRECAUTIONS:

- Store in a refrigerator (2°C-8°C).
- Protect from direct sunlight.
- Do not freeze.
- Keep out of the reach and sight of children.
- Shelf life after first opening: Use the vaccine within 8 hours after first opening.

- The vaccine remains stable for 2 years from production date stated on the label under the recommended storage conditions.

Package sizes:

Bottles of: 50 ml and 100 ml.

FOR VETERINARY USE ONLY.

SPECIAL WARNING(S):

Special warnings for each target species:

Vaccinate healthy animals only.

Special precautions for use in animals:

Not applicable.

Special precautions to be taken by the person administering the medicinal product to animals:

Although Jovaplastm C vaccine is an inactivated vaccine and is not pathogenic for humans; care should be taken to avoid accidental injection. Should accidental injection occur immediate medical care should be sought.

Use during pregnancy, lactation, fertility:

Can be used during pregnancy, lactation and breeding.

Interaction with other medicinal products and other forms of interaction:

No information is available on the safety and efficacy of this vaccine when used with any other veterinary medicinal product. A decision to use this vaccine before and after any other veterinary medicinal product therefore needs to be made on a case by case basis.

Overdose:

2X dose of a vaccine:

Rare: A very small lump at the site of injection may appear in vaccinated animals. This lump usually disappears within 48 hours.

If you notice any side effects please inform your veterinarian.

SPECIAL PRECAUTIONS FOR DISPOSAL:

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal product should be disposed of in accordance with local requirements.

OTHER INFORMATION:

Contagious Caprine Pleuropneumonia (CCPP) is a disease with high morbidity (80-100%) and mortality (60-80%) rates. Classical, acute CCPP is caused by *Mycoplasma capricolum* subspecies *capripneumoniae* (Mccp), originally known as *Mycoplasma* F38 biotype. The disease is characterized by fibrinous Pleuropneumonia with increased straw-colored pleural fluid. The disease is mainly controlled by vaccination.

MANUFACTURING AUTHORIZATION HOLDER AND MANUFACTURED BY:

JORDAN BIO INDUSTRIES CENTER
Tel. +962 6 5232162, Fax: +962 6 5232210
P.O. Box 43, Amman 11941- Jordan
E-mail: info@jovaccnter.com, www.jovaccnter.com



Jovac